

# Confirmatory Factor Models

## *(CFA: Confirmatory Factor Analysis)*

- Topics:
  - **Comparison of EFA and CFA**
  - CFA model parameters
  - Two parts of CFA model identification
  - CFA model estimation
  - CFA model fit evaluation

# EFA vs. CFA: What gets analyzed

- **EFA: Correlation matrix (of items = indicators)**
  - Only correlations among observed item responses are used
  - Only a standardized solution is provided, so the original means and variances of the observed item responses are irrelevant
- **CFA: Covariance matrix (of items = indicators)**
  - Means, variances, and covariances of item responses are analyzed (where the latter two form the “item variance–covariance matrix”)
    - “item covariance matrix” is a shorter way of saying “item variance–covariance matrix”
  - Item response means historically have been ignored (but not by us!)
  - Output includes unstandardized AND standardized solutions
    - **Unstandardized** solution predicts the **item variance–covariance matrix** (regression solution retains original absolute information)
    - **Standardized** (STDYX) solution predicts the **item correlation matrix** (easier to interpret relative sizes of relationships as correlations)

# EFA vs. CFA: Interpretation

- **EFA: Rotation**

- All items load on all **factors** (*aka*, latent traits), no matter what!
- Goal is to pick a rotation that gives closest approximation to “simple structure” (clearly-defined factors, fewest cross-loadings)
- No way of distinguishing latent variables due to “content” (traits being measured) from “method” (correlation created by common approach)
- Summary: **It is NOT your data’s job to tell you what they measure...**

- **CFA: Defining interpretation is your job in the first place!**

- CFA is theory-driven: any structure becomes a testable hypothesis
- You specify number of latent factors and their inter-correlations
- You specify which items load on which latent factors (yes/no)
- You specify any additional relationships for method/other covariance
- You just need a clue; you don’t have to be right (misfit is informative)

# EFA vs. CFA: Judging model fit

- **EFA: Eye-balls and Opinion**

- #Factors? Scree-ish plots, interpretability...
- Which rotation? Whichever makes the most sense... (to you): there is no way to say which is the “right” rotation (and interpretation)!
- Which items load on each factor? Arbitrary cut-off of .3-.4ish
- Standard errors found in newer packages (but still infrequently used)

- **CFA: Inferential tests via Maximum Likelihood (ML or MLR)**

- Global model fit test (and local model fit diagnostics)
- Standard errors (and significance tests) of item loadings, error variances, and error covariances (all parameters)
- Ability to test appropriateness of model constraints or model additions via tests for change in model fit

# EFA vs. CFA: Factor scores

- **EFA: Don't ever use factor scores from an EFA**
  - Factor scores are indeterminate (especially due to rotation)
  - Inconsistency in how factor models are applied to data
    - Factor model is based on common variance only (factors are predictors)
    - Summing items? That's using total variance (components are outcomes)
- **CFA: Factor scores *can* be used, but only if necessary**
  - Best option: Test relations among latent factors directly through SEM
    - Factors can either be predictors ("exogenous" variables) or outcomes ("endogenous" variables) or both at once as needed (e.g., as mediators)
    - Relations between factors will be disattenuated for measurement error
  - Factor scores are less indeterminate in CFA, and could be used
    - In reality, though, factor scores are not known single values because they are modeled as random effects, not fixed effects per person
    - Next-best option: Use "plausible values" or other two-stage approaches that acknowledge uncertainty in factor score estimates (stay tuned)

# Confirmatory Factor Models

## *(CFA: Confirmatory Factor Analysis)*

- Topics:
  - Comparison of EFA and CFA
  - **CFA model parameters**
  - Two parts of CFA model identification
  - CFA model estimation
  - CFA model fit evaluation

# Confirmatory Factor Analysis (CFA)

- **The CFA unit of analysis is the ITEM (as in any LTMM):**

$$y_{is} = \mu_i + \lambda_i F_s + e_{is} \rightarrow \text{both items AND subjects matter}$$

- Observed response for **item  $i$**  and **subject  $s$** 
  - = intercept of item  $i$  ( $\mu$ )
  - + subject  $s$ 's latent factor ( $F$ ), weighted by item-specific loading  $\lambda$
  - + error ( $e$ ) of item  $i$  and subject  $s$

- **This looks like linear regression (with an unobserved X)!**

- $y_s = \beta_0 + \beta_1 X_s + e_s \rightarrow$  written for each item  $\rightarrow y_{is} = \beta_{0i} + \beta_{1i} X_s + e_{is}$
- $\beta_{0i}$  Intercept =  $\mu_i$  = expected item response  $\hat{y}_{is}$  when  $F_s = 0$
- $\beta_{1i}$  Slope of Factor =  $\lambda_i$  = expected difference in  $y_{is}$  for a one-unit difference in  $F_s$  ("difference" because these are  $F$ 's for different people)
- $e_{is}$  Error (Residual) = how far off actual  $y_{is}$  is from predicted  $\hat{y}_{is}$ 
  - Residual has another meaning of "discrepancy" (for model misfit) in latent trait measurement models, so I will try to say "error" for this item parameter instead

# Revisiting Vocabulary: Item Psychometric Properties


- **Item Discrimination**: How related each item is to the latent trait
  - In CTT, discrimination is given by item–total (*aka*, item–remainder) correlation
    - The total score is the best estimate of the latent trait in CTT ( $\rightarrow F_s$  in CFA)
  - In **CFA**, discrimination is given by the **factor loading/slope** ( $\lambda_i$ )
    - We now have a factor that directly accounts for the covariance among items
    - Stronger standardized factor loadings indicate better, more discriminating items
- **Item Difficulty/Severity**: Location of item on the latent trait metric
  - In CTT, difficulty is given by the item mean
  - In **CFA**, difficulty is given by the **item intercept** ( $\mu_i$ ) – which is still backwards
  - In contrast to other latent trait models (such as IFA/IRT), difficulty (intercepts) are often ignored in CFA... here's why...



# Why Item Intercepts Are Often Ignored...

A **“good” item** has a large slope (i.e., factor loading) in predicting the item response from the factor. Because this is a **linear slope**, the item is assumed to be equally discriminating (**equally good**) across the entire latent factor.

Similarly, a **“bad” item** has a flatter linear slope that is **equally bad** across the entire latent factor range (slope=0 means item is unrelated to factor).




Here item intercepts are irrelevant in evaluating how “good” an item is, so they are not really needed in CFA.

But we will estimate them, because item intercepts are critical when:

- Testing factor mean differences in any latent factor model
- Items need to have a nonlinear slope in predicting the item response from the factor (IRT)

# Example Diagram of Two-Factor CFA Model

But some parameters will have to be fixed to known values for the model to be identified.



## Measurement Model for Items:

$\lambda$  = factor loading  
 $e$  = error variance  
 $\mu$  = intercept

## Structural Model for Factors:

$F$  = factor variance  
 $\sigma_{F1,F2}$  = factor covariance  
 $K$  = factor mean

# 2 Types of CFA Parameter Solutions

- Unstandardized → predicts scale-sensitive original item response:
  - **Regression Model:**  $y_{is} = \mu_i + \lambda_i F_s + e_{is}$
  - *Useful when comparing solutions across groups or time (when absolute values matter)*
  - Unstandardized model parameters predict **item means and variance-covariance matrix**
  - Note the solution asymmetry: item parameters  $\mu_i$  and  $\lambda_i$  will be given in the item metric, but  $e_{is}$  will be given as the error variance across persons for that item (squared metric)
  - $Var(y_i) = [\lambda_i^2 * Var(F)] + Var(e_i)$
- Standardized → Solution transformed to  $Var(y_i) = Var(F) = 1$  via **STDYX**:
  - *Useful when comparing items within a solution (relative values on same scale)*
  - Standardized model parameters predict the **item correlation matrix (item means = 0)**
  - Standardized intercept =  $\mu_i / SD(y_{is})$  → not typically reported (and mostly unhelpful)
  - Standardized factor loading =  $[\lambda_i * SD(F_s)] / SD(y_{is}) = \text{item correlation with factor}$
  - Standardized error variance =  $1 - \text{stand } \lambda_i^2 = \text{"item variance due to NOT the factor"}$
  - $R^2$  for item =  $\text{stand } \lambda_i^2 = \text{"item variance due to the factor"}$

Careful! There is also STDY and STD

# Example Two-Factor Model Equations

- Measurement model per numbered item for subject  $s$ :

$$\triangleright y_{1s} = \mu_1 + \lambda_{11}F_{1s} + 0F_{2s} + e_{1s}$$

$$\triangleright y_{2s} = \mu_2 + \lambda_{21}F_{1s} + 0F_{2s} + e_{2s}$$

$$\triangleright y_{3s} = \mu_3 + \lambda_{31}F_{1s} + 0F_{2s} + e_{3s}$$

$$\triangleright y_{4s} = \mu_4 + 0F_{1s} + \lambda_{42}F_{2s} + e_{4s}$$

$$\triangleright y_{5s} = \mu_5 + 0F_{1s} + \lambda_{52}F_{2s} + e_{5s}$$

$$\triangleright y_{6s} = \mu_6 + 0F_{1s} + \lambda_{62}F_{2s} + e_{6s}$$

You decide **how many factors** and if each item has an estimated loading on each factor or not.

Unstandardized loadings ( $\lambda$ ) are **linear slopes** predicting the item response ( $y$ ) from the factor ( $F$ ). **Thus, the model assumes a linear relationship between the factor and the item response.**

**Standardized** loadings are the slopes in a **correlation** metric (and Standardized Loading<sup>2</sup> =  $R^2$ ).

Intercepts ( $\mu_i$ ) are the expected item responses ( $\hat{y}_{is}$ ) when all factors = 0.

Here is the general matrix equation for these 6 item-specific equations:

$$Y = \mu + \lambda F + e$$

where  $Y$ ,  $\mu$ , and  $e$  = 6x1 matrices (because each item gets one value);  $\lambda$  = 6x2 matrix, and  $F$  = 2x1 matrix (because there are two factors)

# The Role of the CFA Model Parameters

- Data going in to be recreated by the CFA model parameters = item means + item variance–covariance matrix
- The CFA **item intercepts** ( $\mu_i$ ) recreate the **item means**
  - Item means are unconditional; item intercepts are conditional on  $F_s = 0$
  - When each item gets its own intercept (the usual case), the item means will be perfectly recreated (**so no room for mis-fit or mis-prediction**)
- The CFA **item error variances** ( $Var[e_i]$ ) recreate the **item variances**
  - Item variances are unconditional; item error variances are conditional (leftover variance after accounting for the contribution of the factor)
  - When each item gets its own error variance (usual case), the item variances will be perfectly recreated (**so no room for misfit or mis-prediction**)
- The CFA **item factor loadings** ( $\lambda_i$ ) recreate the **item covariances**
  - Given 3+ items, there will be more covariances among items to predict than item factor loadings to recreate them, **thus creating room for misfit**

# CFA Model Predictions: ( $F_1$ BY $y_1$ - $y_3$ , $F_2$ BY $y_4$ - $y_6$ )

Items from same factor (room for misfit=bad recreation):

- Unstandardized solution:  $\text{Covariance}(y_1, y_3) = \lambda_{11} * \text{Var}(F_1) * \lambda_{31}$
- Standardized solution:  $\text{Correlation}(y_1, y_3) = \lambda_{11} * (1) * \lambda_{31} \leftarrow \text{std loadings}$
- ONLY reason for correlation is their common factor (local independence, LI)

Items from different factors (room for misfit=bad recreation):

- Unstandardized:  $\text{Covariance}(y_1, y_6) = \lambda_{11} * \text{Cov}(F_1, F_2) * \lambda_{62}$
- Standardized:  $\text{Correlation}(y_1, y_6) = \lambda_{11} * \text{Cor}(F_1, F_2) * \lambda_{62} \leftarrow \text{std loadings}$
- ONLY reason for correlation is the correlation between factors (again, LI)

Variances are additive (and will be reproduced correctly):

- $\text{Var}(y_1) = (\lambda_{11}^2) * \text{Var}(F_1) + \text{Var}(e_i) \rightarrow \text{note imbalance of } \lambda^2 \text{ and } e_i$

# Recreated Item Variance–Covariance Matrix

- Matrix equation:  $\Sigma = \Lambda\Phi\Lambda^T + \Psi$

$\Sigma$  = (Sigma) model-recreated item variance–covariance matrix is built from:

$\Lambda$  = (Lambda) item factor loadings

$\Phi$  = (Phi) factor variances and covariances

$\Lambda^T$  = item factor loadings transposed ( $\rightarrow \lambda^2$ )

$\Psi$  = (Psi) item error (residual) variances

$$\Sigma = \begin{pmatrix} \sigma_{y1}^2 & \sigma_{y1,y2} & \sigma_{y1,y3} & \sigma_{y1,y4} & \sigma_{y1,y5} & \sigma_{y1,y6} \\ \sigma_{y2,y1} & \sigma_{y2}^2 & \sigma_{y2,y3} & \sigma_{y2,y4} & \sigma_{y2,y5} & \sigma_{y2,y6} \\ \sigma_{y3,y1} & \sigma_{y3,y2} & \sigma_{y3}^2 & \sigma_{y3,y4} & \sigma_{y3,y5} & \sigma_{y3,y6} \\ \sigma_{y4,y1} & \sigma_{y4,y2} & \sigma_{y4,y3} & \sigma_{y4}^2 & \sigma_{y4,y5} & \sigma_{y4,y6} \\ \sigma_{y5,y1} & \sigma_{y5,y2} & \sigma_{y5,y3} & \sigma_{y5,y4} & \sigma_{y5}^2 & \sigma_{y5,y6} \\ \sigma_{y6,y1} & \sigma_{y6,y2} & \sigma_{y6,y3} & \sigma_{y6,y4} & \sigma_{y6,y5} & \sigma_{y6}^2 \end{pmatrix} = \begin{pmatrix} \lambda_{11} & 0 \\ \lambda_{21} & 0 \\ \lambda_{31} & 0 \\ 0 & \lambda_{42} \\ 0 & \lambda_{52} \\ 0 & \lambda_{62} \end{pmatrix} \begin{pmatrix} \sigma_{F1}^2 & \sigma_{F1,F2} \\ \sigma_{F2,F1} & \sigma_{F2}^2 \end{pmatrix} \begin{pmatrix} \lambda_{11} & \lambda_{21} & \lambda_{31} & 0 & 0 & 0 \\ 0 & 0 & 0 & \lambda_{42} & \lambda_{52} & \lambda_{62} \end{pmatrix} + \begin{pmatrix} \sigma_{e1}^2 & 0 & 0 & 0 & 0 & 0 \\ 0 & \sigma_{e2}^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sigma_{e3}^2 & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma_{e4}^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sigma_{e5}^2 & 0 \\ 0 & 0 & 0 & 0 & 0 & \sigma_{e6}^2 \end{pmatrix}$$

# Recreated Item Variance–Covariance Matrix

•  $\Sigma = \Lambda\Phi\Lambda^T + \Psi \rightarrow$  **Model-Implied Matrix**

*Sigma for model =*  
 $\Lambda * (\Phi) * \Lambda^T + \Psi$

The **loadings** control how related items from the same factor are predicted to be.

<b>Items within Factor 1</b>					
$\lambda_{11}^2\sigma_{F1}^2 + \sigma_{e1}^2$	$\lambda_{11}\sigma_{F1}\lambda_{21}$	$\lambda_{11}\sigma_{F1}\lambda_{31}$	$\lambda_{11}\sigma_{F2,F1}\lambda_{42}$	$\lambda_{11}\sigma_{F2,F1}\lambda_{52}$	$\lambda_{11}\sigma_{F2,F1}\lambda_{62}$
$\lambda_{21}\sigma_{F1}\lambda_{11}$	$\lambda_{21}^2\sigma_{F1}^2 + \sigma_{e2}^2$	$\lambda_{21}\sigma_{F1}\lambda_{31}$	$\lambda_{21}\sigma_{F2,F1}\lambda_{42}$	$\lambda_{21}\sigma_{F2,F1}\lambda_{52}$	$\lambda_{21}\sigma_{F2,F1}\lambda_{62}$
$\lambda_{31}\sigma_{F1}\lambda_{11}$	$\lambda_{31}\sigma_{F1}\lambda_{21}$	$\lambda_{31}^2\sigma_{F1}^2 + \sigma_{e3}^2$	$\lambda_{31}\sigma_{F2,F1}\lambda_{42}$	$\lambda_{31}\sigma_{F2,F1}\lambda_{52}$	$\lambda_{31}\sigma_{F2,F1}\lambda_{62}$
$\lambda_{42}\sigma_{F2,F1}\lambda_{11}$	$\lambda_{42}\sigma_{F2,F1}\lambda_{21}$	$\lambda_{42}\sigma_{F2,F1}\lambda_{31}$	$\lambda_{42}^2\sigma_{F2}^2 + \sigma_{e4}^2$	$\lambda_{42}\sigma_{F2}^2\lambda_{52}$	$\lambda_{42}\sigma_{F2}^2\lambda_{62}$
$\lambda_{52}\sigma_{F2,F1}\lambda_{11}$	$\lambda_{52}\sigma_{F2,F1}\lambda_{21}$	$\lambda_{52}\sigma_{F2,F1}\lambda_{31}$	$\lambda_{52}\sigma_{F2}^2\lambda_{42}$	$\lambda_{52}^2\sigma_{F2}^2 + \sigma_{e5}^2$	$\lambda_{52}\sigma_{F2}^2\lambda_{62}$
$\lambda_{62}\sigma_{F2,F1}\lambda_{11}$	$\lambda_{62}\sigma_{F2,F1}\lambda_{21}$	$\lambda_{62}\sigma_{F2,F1}\lambda_{31}$	$\lambda_{62}\sigma_{F2}^2\lambda_{42}$	$\lambda_{62}\sigma_{F2}^2\lambda_{52}$	$\lambda_{62}^2\sigma_{F2}^2 + \sigma_{e6}^2$
			<b>Items within Factor 2</b>		


The only reason why items from different factors should be related is the **covariance** between the two factors.

The **loadings** also control how much of the item response is due to factor versus error.



# Example Diagram of Two-Factor CFA Model

But some parameters will have to be fixed to known values for the model to be identified.



## Measurement Model for Items:

$\lambda$  = factor loading  
 $e$  = error variance  
 $\mu$  = intercept

## Structural Model for Factors:

$F$  = factor variance  
 $\sigma_{F1,F2}$  = factor covariance  
 $K$  = factor mean

# Confirmatory Factor Models

## *(CFA: Confirmatory Factor Analysis)*

- Topics:
  - Comparison of EFA and CFA
  - CFA model parameters
  - **Two parts of CFA model identification**
  - CFA model estimation
  - CFA model fit evaluation


# Two Parts of Model Identification

- **Part 1: Create a scale for each latent factor**
  - Each latent factor needs a mean and a variance
  - Necessary but not sufficient for estimating the CFA model
- **Part 2: Ensure the CFA model is estimable**
  - Data going in versus estimated parameters going out:
    - Item means  $\rightarrow$  item intercepts (usually 1:1 ratio)
    - Item variances  $\rightarrow$  item error variances (usually 1:1 ratio)
    - Item covariances  $\rightarrow$  item factor loadings (must have ratio  $\geq 1$ )
  - In practice, this means the number of estimated loadings may not exceed the number of observed item covariances
  - This also means that each pair of variables has only one direct relationship (more of a concern in path models)

# CFA Model Identification Part I:

## *Create a Scale for the Latent Factor Variance*

“Z-Score Factor”  
Method (1)




- The factor doesn't exist, so **it needs a scale** (it needs a mean and variance):
- There are two **equivalent** options to create a scale for the factor **VARIANCE**:

➤ (1) **Fix factor variance to 1: “z-score”**

- Factor SD is same as standard z-score (but any constant  $> 0$  would also be ok)
- **Shouldn't** be used in models with higher-order factors (coming later in this course)

“Marker Item”  
Method (2)





➤ (2) **Fix a “marker item” loading to 1**

- Factor variance is then estimated the “reliable” part of the marker item variance
- e.g., Std. loading = 0.9, item variance = 16? Factor variance =  $(0.9^2) * 16 = 12.96$
- Can cause the model to blow up if marker item has no correlation with the factor at all

# CFA Model Identification Part I:

## Create a Scale for the Latent Factor Mean




- The factor doesn't exist, so **it needs a scale** (it needs a mean and variance):
- There are two **equivalent** options to create a scale for the factor **MEAN**:
  - **(1) Fix factor mean to 0: “z-score”**
    - Factor mean is same as standard z-score (but any constant would also be ok)
    - **Can** be used in models with higher-order factors (coming later in the course)
    - Item intercepts = item means in this case
  - **(2) Fix a “marker item” intercept to 0**
    - Factor mean = mean of marker item
    - Item intercepts = expected item responses when factor = 0 (→ marker = 0, which may not make sense for all item response scales)

# Part I: Possible Factor Means and Variances

## Factor Variance = 1 (fixed)


Factor Mean = 0 (fixed)




## Factor Variance Estimated

Default in Mplus and cfa in lavaan


Factor Mean = 0



Factor Mean Est. = ? (free)



Factor Mean = ?




# Part 2 of CFA Model Identification

- **After scaling the factors**, then try to reproduce observed item means and variance–covariance matrix using as few estimated parameters as possible
  - (Robust) Maximum likelihood used to estimate model parameters
    - **Measurement Model:** Item intercepts, item factor loadings, item error variances
    - **Structural Model:** Factor means, factor variances, factor covariances
  - Global model fit = difference between model-recreated and data-observed variance–covariance matrix (but only covariances usually contribute to misfit)
- How many possible parameters can you estimate: what is total DF?
  - **Total DF** =  $\frac{v(v+1)}{2} + v$  where  $v$  is the # items (NOT people, like usual)
    - Total DF = number of item means, variances, and covariances
    - e.g., if  $v = 4$  items, then Total DF =  $\frac{4(4+1)}{2} + 4 = 14$
  - **Model DF** = data input – model output
  - **Model DF** = # possible parameters – # estimated parameters
  - **Model DF** = # parameter left over (could have been estimated, but didn't)

# Under-Identified Factor: 2 Items

- Model is **under-identified** if there are more unknown parameters than item variances, covariances, and means with which to estimate them
  - Model **cannot be estimated** because there are an infinite number of different parameter estimates that would result in the same (and perfect) fit
  - Example:  $x + y = 7$  ??

You'd have to constrain the loadings to be equal for the model to be estimable.



In other words, the assumptions required to calculate two-score reliability in CTT are the result of model under-identification.

Total possible **DF** = unique pieces of data = **5**

**0 factor variances**

**0 factor means**

**2 item loadings** OR

**2 item intercepts**

**2 error variances**

**1 factor variance**

**1 factor mean**

**1 item loading**

**1 item intercept**

**2 error variances**

**Model DF = 5 - 6 = - 1**

If  $cor(y_1, y_2) = .64$ , then:

$\lambda_{11} = .800, \lambda_{21} = .800$  ??


$\lambda_{11} = .900, \lambda_{21} = .711$  ??

$\lambda_{11} = .750, \lambda_{21} = .853$  ??



# Just-Identified Factor: 3 Items

- Model is **just-identified** if there are as many unknown parameters as item variances, covariances, and means with which to estimate them
  - The model is estimable, so the parameter estimates have a unique solution
  - But parameters will **perfectly reproduce** the observed variance–covariance matrix, so **model fit is not testable**—it's just a re-arrangement of the data
  - Example: Solve  $x + y = 7$ ,  $3x - y = 1$



Total possible **DF** = unique pieces of data = **9**

**0** factor variances

**0** factor means

**3** item loadings

**3** item intercepts

**3** error variances

**1** factor variance

**1** factor mean

**2** item loadings

**2** item intercepts


**3** error variances

OR

**Model DF = 9 - 9 = 0**

Not really a model—more like a description

# Example: Solving a Just-Identified Model




	$y_1$	$y_2$	$y_3$
$y_1$	1.00		
$y_2$	.595	1.00	
$y_3$	.448	.544	1.00

- Step 1:  $ab = .595$   
 $ac = .448$   
 $bc = .544$
- Step 2:  $b = .595/a$   
 $c = .448/a$   
 $(.595/a)(.448/a) = .544$
- Step 3:  $.26656/a^2 = .544$   
 $a = .70$
- Step 4:  $.70b = .595 \quad b = .85$   
 $.70c = .448 \quad c = .64$
- Step 5:  $Var(e_1) = 1 - a^2 = .51$

# Over-Identified Factor: 4+ Items

- Model is **over-identified** if there are fewer unknown parameters than item variances, covariances, and means with which to estimate them
  - The model is estimable, so the parameter estimates have a unique solution
  - But now the parameters will NOT perfectly reproduce the observed matrix  
→ **if Model DF > 0, we can test model fit!**



Total possible **DF** = unique pieces of data = **14**

**0** factor variances

**0** factor means

**4** item loadings

**4** item intercepts

**4** error variances

OR

**1** factor variance

**1** factor mean

**3** item loadings

**3** item intercepts


**4** error variances

$$\text{Model DF} = 14 - 12 = 2$$

**Model fit:** Did we do a “good enough” job reproducing the item covariance matrix with 2 fewer parameters than it was possible to use?

# Oops: Empirical Under-Identification

- Did your model blow up (errors instead of output)? Double-check:
  - Part 1: Make sure each factor has a scale: a mean and a variance
  - Part 2: Make sure you aren't estimating more parameters than you have DF
- Sometimes you can set up your model correctly and it will STILL blow up because of **empirical under-identification**
  - It's not you; **it's your data**—here are two examples of when these models should have been identified, but weren't because of an unexpected 0 relationship



# That Other Kind of Measurement Model...

*(i.e., as used in principal components or partial least squares alternatives to CFA)*

Here is a difference between principal components and factor analysis in terms of “types” of items...

## Factor Model (CFA):


- Composed of “Reflective” or “Effects” items
- Factor is thought to **cause** observed item responses
- Items should be correlated
- **Is identified** with 3+ items (fit is testable with 4+ items)

## Component Model:

- Composed of “Formative” or “Emergent” or “Cause” items
- Component is **result** of observed item responses
- Items may not be correlated
- **Will not be identified** no matter how many items *without additional variables in the model*

# Formative (Component) Models

(see Brown 2015 ch. 8 pp. 322-331)




## Model A Parameters:

- 4 factor loadings/regression paths
  - 1 factor disturbance (variance left over)
  - 10 item correlations
  - 5 item variances
  - 5 item means
- $DF = 20 - 25 = -5$

**Not identified**

Formative measurement models are not identified without including other outcomes or predictors of the formative latent factor



## Model C Parameters:

- 4 factor loadings/regression paths
  - 1 factor disturbance (variance left over)
  - 3 item correlations
  - 5 item variances/error variances
  - 5 item means/intercepts
- $DF = 20 - 18 = 2$

**Identified**

Model C has both formative and reflective indicators—the latter might also be “outcomes”

# Intermediate Summary: CFA

- **CFA is a linear model** in which continuous observed item responses are predicted from latent factors (traits) and error
  - Goal: reproduce observed **item means and variance–covariance matrix** using parameters in measurement model (item intercepts, loadings, error variances) and in structural model (factor means, variances, covariances)
  - Factor model makes specific testable mathematical predictions about how item responses should relate to each other: **loadings predict covariances**
  - Need at least 3 items per latent factor for the model to be identified; need **at least 4 items per latent factor for model fit to be testable**
- CFA framework offers significant advantages over CTT by offering the potential for comparability across samples, groups, and time
  - CTT: No separation of observed item responses from true score
    - Sum across items = true score; item properties belong to that sample only
  - CFA: Latent factor is estimated *separately* from item responses
    - Separates interpretation of person trait levels from specific items given
    - Separates interpretation of item properties from specific persons in sample

# Confirmatory Factor Models


## *(CFA: Confirmatory Factor Analysis)*

- Topics:
  - Comparison of EFA and CFA
  - CFA model parameters
  - Two parts of CFA model identification
  - **CFA model estimation**
  - CFA model fit evaluation



# Where the Answers Come From: The Big Picture of ML Estimation

**ESTIMATOR = Robust Maximum Likelihood;**



**... answers ...**

**Any questions?**

# What all do we have to estimate?


- For example, a model with two correlated factors for  $v = 6$  items:
  - $F_1$  measured by items 1,2,3;  $F_2$  measured by items 4,5,6
  - If we fix both factors to have mean=0 and variance=1, then we need:  
**6 intercepts ( $\mu_i$ ) + 6 factor loadings ( $\lambda_i$ ) + 6 error variances ( $\sigma_{e_i}^2$ ) + 1 factor covariance [ $Cov(F_1, F_2)$ ] = 19 total parameters**
- **Item parameters are FIXED effects** → inference about specific item
  - It's ok if missing data leads to different numbers of total items across persons (assumed *missing at random* when using ML, which is conditionally random)
- What about the all the individual person **factor scores**?
  - The individual factor scores are NOT part of the model—in other words, factor scores are modeled as **RANDOM effects** assumed to be multivariate normal
  - So we need the **factor means, variances, and covariances** as sufficient statistics, but **we don't need** the factor scores for the **individual respondents**

# The End Goals of Maximum Likelihood (ML) Estimation

1. Obtain “most likely” values for each unknown parameter in our model (intercepts, loadings, error variances, factor means, factor variances, factor covariances) → the answers → **the estimates**
2. Obtain some kind of index as to how likely each parameter value actually is (i.e., “really likely” or pretty much just a guess?)  
→ **the standard error (SE) of the estimates (smaller is better)**
3. Obtain some kind of index as to how well the model we’ve specified actually describes the data → **the model fit indices**

**How does all this happen? The magic of multivariate normal...  
(but let’s start with univariate normal first)**

# Univariate Normal Distribution



Univariate Normal PDF:

$$f(y_s) = \frac{1}{\sqrt{2\pi\sigma_e^2}} * \exp\left[-\frac{1}{2} * \frac{(y_s - \hat{y}_s)^2}{\sigma_e^2}\right]$$


Sum over persons for log of  $f(y_i)$  =  
Model Log-Likelihood  $\rightarrow$  Model Fit

- This PDF tells us how **likely** (i.e., **tall**) any value of  $y_s$  is given two things:
  - Conditional mean  $\hat{y}_s$
  - Residual variance  $\sigma_e^2$
- We can see this work using the NORMDIST function in excel!
  - Easiest for **empty** model:  
 $y_s = \beta_0 + e_s$
- We can check our math via software using ML!

# ML via Excel NORMDIST

**Key idea: Normal Distribution formula → data height**

Mean	5.19	5.24	-1.0
Variance	6.56	2.00	-1.5
	<b>Right</b>	<b>Wrong</b>	
Outcome	<b>Log(Height)</b>	<b>Log(Height)</b>	-2.0
1.0	-3.20	-5.76	-2.5
2.1	-2.59	-3.73	-3.0
3.0	-2.22	-2.52	-3.5
4.3	-1.92	-1.49	-4.0
4.6	-1.89	-1.37	-4.5
6.2	-1.94	-1.50	-5.0
7.3	-2.20	-2.33	-5.5
7.6	-2.30	-2.66	-6.0
7.8	-2.38	-2.90	
8.0	-2.46	-3.17	
SUM = Model LL = taller is better			
	<b>-23.09</b>	<b>-27.42</b>	



For a demo in excel, [download this file](#) from PSQF 6270: Generalized Linear Models

# Multivariate Normal for $\mathbf{Y}_s$ :

all  $v = 6$  item responses from person  $s$

$$\text{Univariate Normal PDF: } f(y_s) = (2\pi\sigma_e^2)^{-1/2} * \exp\left[-\frac{1}{2} * (y_s - \mu_s) (\sigma_e^2)^{-1} (y_s - \mu_s)\right]$$

$$\text{Multivariate Normal PDF: } f(\mathbf{Y}_s) = (2\pi)^{-v_s/2} * |\Sigma|^{-1/2} * \exp\left[-\frac{1}{2} * (\mathbf{Y}_s - \boldsymbol{\mu})^T (\boldsymbol{\Sigma}_s)^{-1} (\mathbf{Y}_s - \boldsymbol{\mu})\right]$$

- In our CFA model, the only fixed effects that predict the 6 item responses in  $\mathbf{Y}_s$  are the item intercepts (now  $v = 6$  of them in the vector  $\boldsymbol{\mu}$ )
- CFA model also makes  $\boldsymbol{\Sigma}$ , the **recreated** item variance–covariance matrix:
  - In matrices:  $\boldsymbol{\Lambda}$  = loadings,  $\boldsymbol{\Phi}$  = factor variances and covariances,  $\boldsymbol{\Psi}$  = item error variances
  - Variance of Item  $i$ :  $\text{Var}(y_i) = \lambda_i^2 * \text{Var}(F) + \text{Var}(e_i)$
  - Covariance of items on same factor:  $\text{Cov}(y_1, y_2) = \lambda_{11} * \text{Var}(F_1) * \lambda_{21}$
  - Covariance of items on different factors:  $\text{Cov}(y_1, y_6) = \lambda_{11} * \text{Cov}(F_1, F_2) * \lambda_{62}$
- Uses  $|\boldsymbol{\Sigma}|$  = determinant of  $\boldsymbol{\Sigma}$  = summary of *non-redundant* info
- $\boldsymbol{\Sigma}_s^{-1} \rightarrow$  matrix inverse  $\rightarrow$  like dividing (so can't be 0 or negative)

$$\boldsymbol{\Sigma} = \boldsymbol{\Lambda}\boldsymbol{\Phi}\boldsymbol{\Lambda}^T + \boldsymbol{\Psi}$$

# Now Try Some Possible Answers...

(e.g., for those 19 parameters in this example)

- Plug predictions into **log-likelihood** function, sum over persons:

$$\text{Model (H}_0\text{) Likelihood: } L = \prod_{s=1}^N \left\{ (2\pi)^{-v_s/2} * |\boldsymbol{\Sigma}|^{-1/2} * \exp \left[ -\frac{1}{2} (\mathbf{Y}_s - \boldsymbol{\mu})^T (\boldsymbol{\Sigma}_s)^{-1} (\mathbf{Y}_s - \boldsymbol{\mu}) \right] \right\}$$

$$\text{Model (H}_0\text{) Log Likelihood: } LL = \sum_{s=1}^N \left\{ \left[ -\frac{v_s}{2} \log(2\pi) \right] + \left[ -\frac{1}{2} \log |\boldsymbol{\Sigma}| \right] + \left[ -\frac{1}{2} (\mathbf{Y}_s - \boldsymbol{\mu})^T (\boldsymbol{\Sigma}_s)^{-1} (\mathbf{Y}_s - \boldsymbol{\mu}) \right] \right\}$$

- Try one set of possible parameter values, compute LL (total **height**)
- Try another possible set, compute revised LL....
  - Different algorithms are used to decide which values to try given that each parameter has its own likelihood distribution → like an uncharted mountain
  - Calculus helps the program scale this multidimensional mountain
    - At the top, all first partial derivatives (linear slopes at that point)  $\approx 0$
    - *Positive* first partial derivative? Too *low*, try again. *Negative*? Too *high*.
    - Matrix of partial first derivatives = "score function" = "gradient"

# End Goals 1 and 2: Model Estimates and SEs

- Process terminates (the model “**converges**”) when the next set of tried parameter values don’t improve the LL very much...
  - e.g., Mplus default convergence criteria for this  $H_0$  Model LL = .00005 (other values are used for different estimation problems—see manual)
  - Those are the values for our model parameters that, relative to the other possible values tried, are “most likely” → **Model ( $H_0$ ) LL** and **estimates**
- But we also need to know how trustworthy those estimates are...
  - **Precision** is indexed by the steepness of the multidimensional mountain, where steepness → more negative partial second derivatives
  - Matrix of partial second derivatives = “Hessian matrix”
  - Hessian matrix \*  $-1$  = “information matrix”

Each parameter SE = $\frac{1}{\sqrt{\text{information}}}$
---
  - So steeper function = more information = more precision = **smaller SE**



# End Goal #3: How well do the model predictions match the data?

- Use your model  $LL_{H_0}$  from predicting  $\Sigma \rightarrow$  so how good is it?
- Get the best possible  $LL_{H_1}$  if we used the real data ( $\mathbf{S}$ ) instead:

$$\text{Saturated Model (H}_1\text{) Log Likelihood: } LL = \sum_{s=1}^N \left\{ \left[ -\frac{v_s}{2} \log(2\pi) \right] + \left[ -\frac{1}{2} \log|\mathbf{S}| \right] + \left[ -\frac{1}{2} v_s \right] \right\}$$

- Compute the **ML fitting function** that indexes how far off the model recreations are from the real data moments  $\rightarrow \chi^2$ :

$$\text{ML Fitting Function: } F_{ML} = \frac{LL_{H_1 \text{ data}}}{N} - \frac{LL_{H_0 \text{ model}}}{N} \quad \text{where} \quad \chi^2 = 2 * N * F_{ML}$$

- Combining and re-arranging the terms in LL for  $H_0$  and  $H_1$  yields this common (complete data) expression for the ML fitting function:

$$F_{ML} = \frac{1}{2} \sum_{s=1}^N \left\{ \underbrace{\log|\Sigma| - \log|\mathbf{S}|}_{\text{how far off}} + \underbrace{\text{trace}[(\Sigma)^{-1} \mathbf{S}] - v_s}_{\text{correction for \#items}} \right\} / N$$

If the model fits perfectly, both parts should be 0.

# What about item non-normality?

- The use of this multivariate normal ML function assumes:
  - Persons and items are conditionally independent
  - Item responses can be missing at random (MAR; ignorable)
  - Factor scores ( $F_s$ ) have a multivariate normal distribution
  - Item residuals ( $e_{is}$ ) have a multivariate normal distribution
  - So in this case, the original item responses should have a multivariate normal distribution, too (given prediction by normal  $F_s$  + normal  $e_{is}$ )
- Impact of non-normality of item responses:
  - Linear model predicting item response from factor may not work well
    - if  $y_{is}$  is not really continuous, the slope needs to shut off at its boundaries
  - SEs and  $\chi^2$ -based model fit statistics will be incorrect
  - Three fixes: **1. Robust ML** (or 2. transform the data, or 3. use a different kind of factor model → IRT/IFA... stay tuned)

# Robust ML for Non-Normality: MLR

- **MLR in Mplus:**  $\approx$  Yuan-Bentler  $T_2$  (permits MCAR or MAR missing data)
  - Still a **linear model** between the item responses and latent factor, so the parameter estimates will be the same as in regular ML
- Adjusts **fit statistics** using an estimated **scaling factor**  $\rightarrow$  for kurtosis:
  - Scaling factor = 1.000 = perfectly multivariate normal  $\rightarrow$  same as regular ML!
  - Scaling factor  $>$  1.000 = leptokurtosis (too-fat tails; fixes too-big  $\chi^2$ )
  - Scaling factor  $<$  1.000 = platykurtosis (too-thin tails; fixes too-small  $\chi^2$ )
- **SEs** computed with Huber-White “sandwich” estimator  $\rightarrow$  uses an information matrix from the variance of the partial first derivatives to correct the information matrix from the partial second derivatives
  - Leptokurtosis (too-fat tails)  $\rightarrow$  increases information; fixes too small SEs
  - Platykurtosis (too-thin tails)  $\rightarrow$  lowers information; fixes too big SEs
  - See Enders (2010 ch. 5) for a readable explanation of how MLR differs from ML
- Because MLR simplifies to ML if the item responses actually are multivariate normally distributed, **we will use MLR as our default estimator for CFA**

# SEM Estimation in STATA v. 16

- Although SEMs can be estimated using STATA, it appears there are (currently) fewer combinations allowed than in Mplus:
  - SEM option `method()` allows estimator choices:
    - ML = regular limited-information ML (no missing data allowed)
    - MLMV = full-information ML (MCAR or MAR missing data)
    - ADF = asymptotic distribution free (requires huge  $N$  for stable estimation)
  - SEM option `vce()` allows robust standard error choices:
    - Robust = Huber-White “sandwich” version (as given by MLR in Mplus)
      - Can be used with MLMV for missing data to adjust parameter SEs
      - No scaling correction factor given to compute fit statistics (none given)
    - Sbentler = Satorra–Bentler version (MLM in Mplus)
      - Can only be used with ML estimation method (so no missing data allowed)
- Because there appears to be no combination that allows missing data + robust estimation of fit statistics and SEs, I’m not going to provide STATA SEM example code...

# Confirmatory Factor Models

## *(CFA: Confirmatory Factor Analysis)*


- Topics:
  - Comparison of EFA and CFA
  - CFA model parameters
  - Two parts of CFA model identification
  - CFA model estimation
  - **CFA model fit evaluation**

# The Big Picture of Model Fit

- Aspects of the observed data to be predicted  
(assuming a z-score metric for the factors for simplicity):
- CFA model equation:  $\mathbf{y}_{is} = \boldsymbol{\mu}_i + \boldsymbol{\lambda}_i \mathbf{F}_s + \mathbf{e}_{is}$ 
  - **Mean** per item: **Captured** by intercept  $\boldsymbol{\mu}_i$  per item
    - Not a source of misfit (unless constraints are applied on the intercepts)
  - **Variance** per item: **Captured** by weighted factor + error
    - $\text{Var}(\mathbf{y}_i) = \boldsymbol{\lambda}_i^2 * \text{Var}(\mathbf{F}) + \text{Var}(\mathbf{e}_i) \rightarrow$  output given as  $\boldsymbol{\lambda}_i$  and  $\text{Var}(\mathbf{e}_i)$
    - Factor and error variances are additive  $\rightarrow$  not a source of misfit  
(whatever  $\mathbf{F}_s$  doesn't get,  $\mathbf{e}_{is}$  picks up to get back to total  $\mathbf{y}_i$  variance)
  - **Covariance** among items: **Predicted** via factor loadings  $\boldsymbol{\lambda}_i$ 
    - Loadings (multiplied) predict what observed covariance should be...  
but they may not be right  $\rightarrow$  **THE PRIMARY SOURCE OF MISFIT**

# Baselines for Assessing Fit in CFA

(Item means all saturated in both)



# Baseline model comparisons are already given in Mplus output (MLR here)

## MODEL FIT INFORMATION (Abbreviated)

# Free parameters = # estimated

Number of Free Parameters

18

### Loglikelihood

H0 Value -11536.404  
 H0 Scaling Correction Factor 1.4158  
 for MLR  
 H1 Value -11322.435  
 H1 Scaling Correction Factor 1.4073  
 for MLR

$H_1$  Saturated (Unstructured) Model Height from this  $\rightarrow$

$$\begin{pmatrix} \sigma_{y1}^2 & \sigma_{y1,y2} & \sigma_{y1,y3} & \sigma_{y1,y4} & \sigma_{y1,y5} & \sigma_{y1,y6} \\ \sigma_{y2,y1} & \sigma_{y2}^2 & \sigma_{y2,y3} & \sigma_{y2,y4} & \sigma_{y2,y5} & \sigma_{y2,y6} \\ \sigma_{y3,y1} & \sigma_{y3,y2} & \sigma_{y3}^2 & \sigma_{y3,y4} & \sigma_{y3,y5} & \sigma_{y3,y6} \\ \sigma_{y4,y1} & \sigma_{y4,y2} & \sigma_{y4,y3} & \sigma_{y4}^2 & \sigma_{y4,y5} & \sigma_{y4,y6} \\ \sigma_{y5,y1} & \sigma_{y5,y2} & \sigma_{y5,y3} & \sigma_{y5,y4} & \sigma_{y5}^2 & \sigma_{y5,y6} \\ \sigma_{y6,y1} & \sigma_{y6,y2} & \sigma_{y6,y3} & \sigma_{y6,y4} & \sigma_{y6,y5} & \sigma_{y6}^2 \end{pmatrix}$$

"Model fit"  $\chi^2$  is from a  $-2\Delta LL$  test of your chosen  $H_0$  model vs. saturated  $H_1$  model

### Chi-Square Test of Model Fit

Value 307.799\*  
 Degrees of Freedom 9  
 P-Value 0.0000  
 Scaling Correction Factor 1.3903  
 for MLR

### Independence (Null) Model

"Baseline model"  $\chi^2$  is from  $-2\Delta LL$  test of null model vs. saturated  $H_1$  model (ignore)

$$\begin{pmatrix} \sigma_{y1}^2 & 0 & 0 & 0 & 0 & 0 \\ 0 & \sigma_{y2}^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sigma_{y3}^2 & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma_{y4}^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sigma_{y5}^2 & 0 \\ 0 & 0 & 0 & 0 & 0 & \sigma_{y6}^2 \end{pmatrix}$$

### Chi-Square Test of Model Fit for the Baseline Model

Value 1128.693  
 Degrees of Freedom 15  
 P-Value 0.0000



# Baseline model comparisons are already given in R Lavaan output (ML & MLR)

Estimator	ML	Robust
Minimum Function Test Statistic	427.937	307.804
Degrees of freedom	9	9
P-value (Chi-square)	0.000	0.000
Scaling correction factor for the Yuan-Bentler correction (Mplus variant)		1.390

This version of "minimum function test statistic" is  $\chi^2$  from  $-2\Delta LL$  test of your chosen  $H_0$  model vs. saturated  $H_1$  model

Model test baseline model:		
Minimum Function Test Statistic	1981.034	1128.693
Degrees of freedom	15	15
P-value	0.000	0.000

## Independence (Null) Model

"Model test baseline model" version is  $\chi^2$  from  $-2\Delta LL$  test of null model vs. saturated  $H_1$  model (ignore)

$$\begin{pmatrix} \sigma_{y1}^2 & 0 & 0 & 0 & 0 & 0 \\ 0 & \sigma_{y2}^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & \sigma_{y3}^2 & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma_{y4}^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sigma_{y5}^2 & 0 \\ 0 & 0 & 0 & 0 & 0 & \sigma_{y6}^2 \end{pmatrix}$$

[other fit indices omitted for now]

Loglikelihood and Information Criteria:		
Loglikelihood user model (H0)	-11536.404	-11536.404
Scaling correction factor for the MLR correction		1.416
Loglikelihood unrestricted model (H1)	-11322.435	-11322.435
Scaling correction factor for the MLR correction		1.407

Number of free parameters	18	18
Akaike (AIC)	23108.808	23108.808
Bayesian (BIC)	23198.912	23198.912
Sample-size adjusted Bayesian (BIC)	23141.739	23141.739

$H_1$  Saturated (Unstructured) Model Height from this  $\rightarrow$

$$\begin{pmatrix} \sigma_{y1}^2 & \sigma_{y1,y2} & \sigma_{y1,y3} & \sigma_{y1,y4} & \sigma_{y1,y5} & \sigma_{y1,y6} \\ \sigma_{y2,y1} & \sigma_{y2}^2 & \sigma_{y2,y3} & \sigma_{y2,y4} & \sigma_{y2,y5} & \sigma_{y2,y6} \\ \sigma_{y3,y1} & \sigma_{y3,y2} & \sigma_{y3}^2 & \sigma_{y3,y4} & \sigma_{y3,y5} & \sigma_{y3,y6} \\ \sigma_{y4,y1} & \sigma_{y4,y2} & \sigma_{y4,y3} & \sigma_{y4}^2 & \sigma_{y4,y5} & \sigma_{y4,y6} \\ \sigma_{y5,y1} & \sigma_{y5,y2} & \sigma_{y5,y3} & \sigma_{y5,y4} & \sigma_{y5}^2 & \sigma_{y5,y6} \\ \sigma_{y6,y1} & \sigma_{y6,y2} & \sigma_{y6,y3} & \sigma_{y6,y4} & \sigma_{y6,y5} & \sigma_{y6}^2 \end{pmatrix}$$

# Free parameters = # estimated

# 4 Steps in Assessing Model Fit

## 1. Global model fit

- *Does the model “work” overall: Does it adequately recreate the observed item means, variances, and covariances?*
- Recent research has indicated that “standard” cut-offs by which to judge global model fit may not work for all data analyses...

## 2. Local model fit

- *Are there any more specific problems (that cause global misfit)?*

## 3. Inspection of model parameters

- *Are the estimates, SEs, and the predicted item responses plausible?*

## 4. Reliability and information per item

- *How “good” is my test? How useful is each item?*

# Step 1: Indices of Global Model Fit

- Primary fit index: obtained model  $\chi^2 = 2 * N * F_{ML}$ 
  - $\chi^2$  is evaluated based on model DF (# parameters left over)
  - Tests null hypothesis that  $\Sigma = S$  (that model = data is perfect), so **significance is bad** (i.e., smaller  $\chi^2$ , bigger  $p$ -value is better)
    - Is LRT ( $-2\Delta LL$ ) of your  $H_0$  model versus saturated best  $H_1$  model
    - Btw, don't use "ratio rules" like  $\chi^2/DF > 2$  or  $\chi^2/DF > 3$
  - Just using  $\chi^2$  to index model fit is usually insufficient, however:
    - $\chi^2$  depends largely on sample size (is overpowered with large  $N$ )
    - Is "unreasonable" null hypothesis (perfect fit, really??)
    - Btw,  $\chi^2$  fit test is only possible given balanced data (as typical for CFA)
- Because of these issues, additional fit indices are usually used in conjunction with the  $\chi^2$  test (that function like effect sizes)
  - **Absolute** Fit Indices (besides  $\chi^2$ )—relative to "**saturated**" best model
  - **Comparative** (Incremental) Fit Indices—relative to "**null**" worst model
  - Cite a reference for any cut-offs you use... it's now more complicated!

# Step 1: Indices of Global Model Fit

- Absolute Fit: **SRMR**

- **Standardized Root Mean Square Residual**

- Get difference of standardized  $\mathbf{S} - \Sigma \rightarrow$  "residual" (discrepancy) matrix

- Sum the squared residuals of the predicted correlation matrix across items, divide by number of matrix elements, then take square root:

- $$SRMR = \sqrt{\frac{2 \sum_{i=1}^I \sum_{j=1}^J \left[ \frac{s_{ij} - \sigma_{ij}}{s_{ii} s_{jj}} \right]^2}{I(I-1)}}$$

- Ranges from 0 to 1: **smaller is better**

- Convention: ".08 or less"  $\rightarrow$  good fit

- Less common variant: **RMR (Root Mean Square Residual)**

# Step 1: Indices of Global Model Fit

## Parsimony-Corrected: **RMSEA**

- **Root Mean Square Error of Approximation**
- Relies on a “non-centrality parameter” (NCP) for  $T$  (target  $H_0$ )
  - NCP indexes how far off your model is → adjusted  $\chi^2$  distribution
  - $NCP_T = \max(\chi_T^2 - DF_T, 0)$  → scaled discrepancy  $d_T = NCP_T/N$
  - $RMSEA = \sqrt{\frac{\max(\chi_T^2 - DF_T, 0)}{DF_T * N}} = \sqrt{\frac{d}{DF_T}}$  → how far off per Model DF left
- RMSEA ranges from 0 to 1; **smaller is better**
  - Conventions:  $< .05$  or  $.06$  = “good”,  $.05$  to  $.08$  = “adequate”
  - In addition to point estimate, get 90% confidence interval (CI)
  - RMSEA penalizes for model complexity—it’s discrepancy in fit per Model DF left (but not sensitive to  $N$ , although CI can be)
  - Also get test of “close fit”: null hypothesis that  $RMSEA \leq .05$

# Step 1: Indices of Global Model Fit

## Comparative (Incremental) Fit Indices (bigger is better)

- Fit evaluated relative to “null” (independence) model of 0 covariances
- Relative to that terrible baseline, your model fit should be great!
- Conventions:  $> .90$  = “adequate”,  $> .95$  = “good”

## • CFI: Comparative Fit Index (ranges from 0 to 1)

➤ Also based on idea of NCP ( $\chi_T^2 - DF_T$ )

➤ 
$$CFI = \frac{\max(\chi_N^2 - DF_N, 0) - \max(\chi_T^2 - DF_T, 0)}{\max(\chi_N^2 - DF_N, 0)}$$

$T$ = target model ( $H_0$ ) $N$ = null model (no covariances)
---

## • TLI: Tucker-Lewis Index (= Non-Normed Fit Index)

➤ 
$$TLI = \frac{\frac{\chi_N^2}{DF_N} - \frac{\chi_T^2}{DF_T}}{\frac{\chi_N^2}{DF_N} - 1}$$
 (so can go negative or  $> 1$ )

# 4 Steps in Model Evaluation


## 1. Assess global model fit (summary)

- Recall that item intercepts → item means, and item error variances → item variances, so *misfit comes from badly recreated covariances*
- $\chi^2$  is sensitive to large  $N$ , so pick at least one global fit index from each class; hope they agree (e.g., CFI, RMSEA) that fit is “good”
- Conventions of “good” absolute model fit largely stem from simulation studies reported in [Hu & Bentler \(1999\)](#)
  - Been cited 94,000+ times! But no one study can cover everything...
    - Held indicator reliability relatively constant: standardized loadings .70-.80
    - Small-ish model of 15 indicators measuring 3 correlated factors
    - Complete data, generated using perfectly multivariate normal indicators
  - Research now suggests standards for what is “good” model fit will vary significantly as a function of these unaddressed features...
    - Here are examples from recent studies (on your reading list or reference given)


# Good Fit is Easier with Missing Data

- [Zhang & Savalei \(2020\)](#): Cases that don't have responses to the indicators with mis-specification will contribute better fit (higher  $LL$ )!
  - Figure 4: Fit when a correlated residual (error) of increasing size is ignored

RMSEA gets less worse with more mis-specification when missing the indicators that have ignored correlated residuals



CFI also gets less worse with more mis-specification when missing the indicators that have ignored correlated residuals







# Good Fit is Easier with Missing Data

- [Zhang & Savalei \(2020\)](#): Same problem when misfit is due to structural mis-specification (i.e., not localized to indicator errors)
  - Figure 5: Fit when one factor is specified instead of two correlated factors

RMSEA gets less worse with more mis-specification with greater amounts of missing indicators with more relevance

CFI also gets less worse with more mis-specification with greater amounts of missing indicators with more relevance





# Good Fit by Number of Indicators...

- ...It's complicated... see [Shi, Lee, & Maydeu-Olivares \(2019\)](#)
  - Figures 1 and 3: Effects of # indicators for  $N=200, 500, 1000$ , and population

CFI gets more worse with more indicators, smaller  $N$ , and low reliability (for  $\lambda = .40$ , CFI is much more variable)


Correct model: RMSEA gets a little worse (still ok) with more indicators and smaller  $N$   
 Incorrect models: RMSEA gets better with more indicators (less so with small  $N$ )



Right: Mis-specified residuals (errors)  $\rightarrow$  misfit limited to only some indicators (so having more properly specified indicators makes fit better on average)

# Good Fit\* Is Easier With Lower Reliability


- Lower reliability  $\rightarrow$  Lower standardized factor loadings
  - [McNeish, An, & Hancock \(2018\)](#): 15 indicators measuring 3 factors
  - Figures 2 and 3: missing factor covariance  $\rightarrow$  always good fit if  $\lambda = .40!$
  - Strong signal (i.e., more reliability) makes it easier to detect when model does not adequately capture that signal



Wide variability in CFI with  $\lambda = .40!$

# When Fit Indices Disagree

- \*Opposite pattern also found for CFI using more incorrect models: CFI was lower (worse fit) with lower reliability ([Heene et al. 2011](#))
- When might RMSEA and CFI disagree? It's a complex function of amount of misfit and DF with which to test it (as well as reliability)



- Figure 1 from [Lai & Green \(2016\)](#)
  - x-axis = amount of misfit in your  $H_0$  model (up to null model,  $F_b$ )
  - y-axis = model fit discrepancy function;  $>0$  = CFI happier,  $<0$  = RMSEA happier
- In summary: How to know what cutoffs are reasonable can be tricky... see [West et al. \(in press\)](#) for a review (on your reading list)

# 4 Steps in Model Evaluation

## 1. Assess global model fit (summary)

- Recall that *misfit generally comes from badly recreated covariances*
  - Be aware that artificially good absolute fit can be created by indicators with low reliability and/or missing data; assessments of global fit can be more variable with smaller  $N$  in large models
  - Corrections for non-normality also continually being developed...
- If model fit is not good (yet), you should NOT interpret the model parameters, because they will change as the model specification changes
    - If model fit is not good, you need to find out WHY → go to step 2
  - Even if model fit IS good, it does not mean you are done: still proceed to step 2, assessing local fit
    - Good global model fit means that the observed and recreated item means, variances, and covariances aren't too far off on the whole... this doesn't guarantee each specific covariance is recreated well

# 4 Steps in Model Evaluation: Step 2

## 2. Identify local misfit: localized model strain

- What is the discrepancy (“residual”) between each model-recreated and data-observed item mean, variance, and covariance?
- Should inspect **normalized model residuals** for that
  - **RESIDUAL** output option in Mplus, residual () in R lavaan, or ESTAT RESIDUAL in STATA
  - “Normalized” is residual/SE → **works like a z-score**
  - Relatively large absolute values indicate “localized strain” (misfit)
  - **Positive** residual → Items are **more** related than you predicted
    - More than just the factor (your model) creating a covariance
  - **Negative** residual → Items are **less** related than you predicted
    - Not as related as your model said they should be
- **Evidence of localized strain tells you where the problems are, but not how to fix them... (positive is easier than negative)**

# 4 Steps in Model Evaluation: Step 2

2. Identify localized model strain, continued...
  - Parallel info: **Modification Indices** (*aka*, cheat codes)
    - LaGrange Multiplier: decrease in model fit  $\chi^2$  by adding the listed model parameter (e.g., cross-loading, error covariance)
      - Usually only pay attention if  $\chi^2 > 3.84$  for  $DF = 1$  (for  $p < .05$ )
      - Get expected parameter estimate for what's to be added, but should only pay attention if its effect size is "meaningful"
      - Also only pay attention if you can INTERPRET AND DEFEND IT
    - Implement these ONE AT A TIME, because one addition to the model can alter the rest of the model substantially
  - Keep in mind that these "manipulation indices" can only try to repair your current model; **they will never suggest a new factor structure instead** (that's your job)!

# Testing Fixes to the Model

- Most common approach for assessing whether adding or subtracting parameters changes model fit is the likelihood ratio test (aka,  $-2\Delta LL$  “deviance difference” test)
  - Done for you in two cases: comparing saturated  $H_1$  to your  $H_0$  as model  $\chi^2$ , and comparing saturated  $H_1$  to “null” model
  - Implemented via direct difference in model  $\chi^2$  values most often, but this is only appropriate when using regular ML estimation
- Variants of ML for non-normal data (like MLR) require a modified version of this  $-2\Delta LL$  test (see Mplus website):  
<http://www.statmodel.com/chidiff.shtml>
  - Is called “rescaled likelihood ratio test” because it has extra steps to incorporate the scaling factors for each model to be compared
  - I built you a spreadsheet for this when using Mplus...you’re welcome 😊
  - Can also use `anova()` in R lavaan (not sure about LRTEST in STATA)



# Testing Fixes to the Model: $-2\Delta LL$

- Comparing nested models via a “**likelihood ratio test**” →  $-2\Delta LL$  (MLR rescaled version)

Note: Your LL will always be listed as the **H0** (H1 is for the saturated, perfectly fitting model)

➤ 1. Calculate  $-2\Delta LL = -2*(LL_{\text{fewer}} - LL_{\text{more}})$

➤ 2. Calculate **scaling correction for difference** =

$$\frac{(\#parms_{\text{fewer}} * scale_{\text{fewer}}) - (\#parms_{\text{more}} * scale_{\text{more}})}{(\#parms_{\text{fewer}} - \#parms_{\text{more}})}$$

Fewer = simpler model  
More = more parameters

➤ 3. Calculate **rescaled difference** =  $-2\Delta LL / \text{scaling correction}$

➤ 4. Calculate  **$\Delta df$**  =  $\#parms_{\text{more}} - \#parms_{\text{fewer}}$

➤ 5. **Compare rescaled difference to  $\chi^2$  with  $df = \Delta df$**

- Add 1 parameter?  $LL_{\text{diff}} > 3.84$ , add 2 parameters:  $LL_{\text{diff}} > 5.99...$
- Absolute values of LL are meaningless (is relative fit only)
- Process generalizes to many other kinds of models (scale factor of 1 = regular ML)

# Testing Fixes to the Model: $-2\Delta LL$

- If **adding** a parameter, model fit can either get **better** OR stay the same ("not better"):
  - Better = larger LL for  $H_0$  and smaller model  $\chi^2$
  - e.g., add another factor, add error covariance
- If **removing** a parameter, model fit can either get **worse** OR stay the same ("not worse")
  - Worse = smaller LL for  $H_0$  and larger model  $\chi^2$
  - e.g., constrain item loadings equal  $\rightarrow$  test "tau-equivalence"
- When testing parameters that have a boundary (e.g., factor correlation  $\neq 1$ ?), this test will be slightly conservative
  - Should use  $p < .10$  instead of  $p < .05$  (or mixture  $\chi^2$  distribution)
  - Same problem as when testing new random effect variances in MLM


# Testing Fixes to the Model, cont.

- For comparing **non-nested models** (e.g., should  $y_1$  load on  $F_2$  or  $F_1$  instead?), the  $-2\Delta LL$  test is not applicable given same DF
- Use information criteria instead: **AIC** and **BIC**
  - Akaike IC:  $AIC = -2LL + 2*\#\text{parameters}$
  - Bayesian (Schwartz) IC =  $-2LL + \log(N)*\#\text{parameters}$
  - Are NOT significance tests, just “smaller is better”, is “evidence”
  - **Still cannot be used on models with different items (outcomes)**
- For both nested or non-nested model comparisons, differences in other fit indices should be examined, too
  - No real critical values for changes in other fit indices, however
  - They may disagree (especially RMSEA, which likes parsimony)


# Fixing the Model by Expanding

- A common (and relatively easy to fix) source of misfit is caused by items that are too correlated after accounting for their common factor—some possible solutions:
  - Add **error covariance(s)** (i.e., as suggested by cheat code mod indices)
    - Is additive:  $Cov(y_1, y_2) = \text{cov due to Factor} + \text{cov due to error covariance}$ , so the residual covariance basically plugs the hole in the covariance matrix
    - In models that do not allow error covariances (e.g., IFA, stay tuned), you can do the same via a separate uncorrelated “**method factor**” (for positive covariance, fix both loadings = 1; for negative covariance, use 1 and -1)
    - **Either way, this means you have unaccounted for multidimensionality**  
→ Explicit acknowledgement that you have measured your latent factor + something else that those items have in common (e.g., stem, valence, specific content) of unknown origin, so you must be able to defend error covariances
  - Lots of problematic pairings? **Re-consider factor dimensionality**
    - I'd generally recommend against adding cross-loadings, because if the item measures more than one thing, it will complicate the interpretation of factors


# Equivalent Ways of Addressing Multidimensionality... (Brown, 2015 p. 181)




Error Covariance



Factor + Method Factor



Two-Factor Models



Here a general factor of "Social Interaction Anxiety" includes two items about public speaking specifically. The extra relationship between the two public speaking items can be modeled in different, yet statistically equivalent ways... **error covariances represent another factor** (which is why you should be able to explain and predict them if you include them).

# When to Simplify the Model

- Factors correlated  $> .85$ ish may suggest a simpler structure
  - Nested model comparison: fix factor variances to 1 so factor covariance becomes factor correlation, then test  $r \neq 1$  at  $p < .10$  (because  $r$  is bounded from  $-1$  to  $1$ )
- When might you consider dropping an item?
  - Non-significant loadings: If the item isn't related, it is NOT measuring the latent trait, and so you probably don't need it
  - Negative loadings: Make sure to reverse-coded as needed ahead of time, otherwise, this indicates a big problem!
  - Problematic leftover positive covariances between two items—such redundancy implies you may not need both (redundancy may indicate a “bloated specific”)
  - If one item is responsible for many of the suggested error covariances, perhaps you might remove it (but be cautious, because often fewer items  $\rightarrow$  less reliability)
- **However: models with different items (outcomes) are NOT COMPARABLE AT ALL because their LL values are based on different input data!**
  - No model comparisons of any kind (including  $-2LL$ , AIC, and BIC)
  - To do a true comparison, you'd need to leave the item in the model but set its loading = 0 (which is the same as the original test of its loading)

# What else can go wrong?

- Error message: “**non-positive definite (NPD)**”
  - Both  $\mathbf{S}$  (data) and  $\mathbf{\Sigma}$  (recreated) matrices must be positive definite
    - Because they get inverted in the LL formula (like matrix division)
  - Non-positive definite means that the determinant is  $\approx 0$ , or that the matrix is singular (has redundant information)
    - Double-check that data are being read in correctly; otherwise you may need to drop items that are too highly correlated
    - **NPD means your model is broken and you can't keep it**
- **Structural under-identification**
  - Does every factor have a mean and variance and at least 3 items?
  - Does the marker item actually load on the factor???
- **Empirical under-identification**
  - More likely with smaller sample sizes, fewer indicators per factor, and items with low reliabilities ( $R^2$  accounted for by factor)

# Open in case of emergency...

- If good model fit seems hopeless, you may choose to go back to the exploratory drawing board... almost
  - Actual EFA uses weird constraints to identify the model, so don't use it
- Brown (2015) suggests an "E/CFA" approach of estimating an exploratory-like model staying within a CFA framework:
  - Fix each factor variance to 1 and mean to 0 for identification
  - Each factor gets one item that loads ONLY on it (loading fixed to 1)
  - Rest of items can load on all factors (or those most likely measured)
  - Why bother? To get significance tests of factor loadings
  - May suggest a useful alternative structure, which should then ideally be replicated in an independent sample using CFA
  - Ideally, any model modifications should be replicated in a new sample!



# Summary: Model Evaluation Steps 1 and 2

## 1. Assess global model fit

- Recall that item intercepts, factor means, and variances are usually just-identified → *so misfit comes from badly recreated covariances*
- $\chi^2$  is sensitive to large sample size, so pick at least one global fit index from each class (e.g., CFI, RMSEA); use cutoffs with caveats

## 2. Identify localized model strain

- Good global model fit means that the observed and recreated item means, variances, and covariances aren't too far off on the whole... this doesn't guarantee each specific covariance is recreated well
- Consider normalized residuals and modification indices to try and "fix" your model (add or remove factors, add or remove residual covariances, etc.)—Has to be theoretically justifiable!!

**Good global and local fit? Great, but we're not done yet...**

# 4 Steps in Model Evaluation: Step 3

## 3. Inspect **parameter effect sizes** and significance

- A 1-factor model will fit each of these correlation matrices perfectly:

	y1	y2	y3	y4
y1	1			
y2	.1	1		
y3	.1	.1	1	
y4	.1	.1	.1	1

	y1	y2	y3	y4
y1	1			
y2	.8	1		
y3	.8	.8	1	
y4	.8	.8	.8	1


- **Good model fit does not guarantee a good model**
- **A good model has meaningful factor loadings**
- **If your items are not correlated, game over, regardless of fit**

# 4 Steps in Model Evaluation: Step 3

3. Inspect **parameter effect sizes** and significance
  - Model fit does not guarantee meaningful factor loadings
    - Can reproduce lack of covariance quite well and still not have anything useful—e.g., factor loading of 0.2 → 4% shared variance?!?
    - **Effect size ( $R^2$  of item variance from factor) is practical significance**
  - Get SEs and  $p$ -values for unstandardized estimates (at least report estimate from standardized solution)
    - Marker items won't have significance tests for their unstandardized loadings because they are fixed at 1, but you'll still get standardized factor loadings for them (help to judge relative importance)
  - Make sure all estimates are within bounds AND predicted item responses are plausible at expected latent factor values ( $\pm 2$  SD)
    - No standardized factor loadings  $> 1$  (unless the indicator has cross-loadings, in which case this is actually possible)
    - No negative factor variances or negative error variances

# 4 Steps in Model Evaluation: Step 3

- CFA is a linear model, so you can plot the responses predicted from the unstandardized item intercepts and loadings (slopes) across factor values
- If predicted responses exceed possible range within  $\pm 2$  SD of factors, then a **linear CFA may not be appropriate**




- CFA using logit-transformed item responses is a potential solution for bounded/skewed continuous items (creates a logistic curve)

➤  $L = \min - 1, U = \max + 1, \text{Logit} = \text{LOG} \left( \frac{y_{is} - L}{U - y_{is}} \right)$

➤ Predicted  $y_{is} = L + (U - L) \frac{\exp(\text{Logit})}{1 + \exp(\text{Logit})}$

- Alternatively, [CFA using a beta distribution!](#)
- For ordinal responses, an IFA/IRT model is a better option (stay tuned)



# 4 Steps in Model Evaluation: Step 4

## 4. Calculate item information and model-based reliability

- **Item Information** =  $(\text{unstandardized } \lambda)^2 / \text{Var}(e)$ 
  - What proportion of item variance is “true” relative to error?
    - Size of unstandardized loadings by themselves is not enough, as their relative contribution depends on size of error variance
    - The **standardized loadings** will give you the same rank order in terms of item information, which is why information is not often used within CFA (but stay tuned for item and test information in IRT/IFA models)

- **“Omega” Sum Score Reliability** = 
$$\frac{\text{Var}(F) * (\Sigma \lambda)^2}{[\text{Var}(F) * (\Sigma \lambda)^2] + \Sigma \text{Var}(e) + 2 \Sigma (e \text{ cov})}$$

- Factor variance \* squared sum of unstandardized factor loadings, over that + summed error variances + 2\*summed error covariances
  - Although Omega should be calculated using unstandardized loadings, Omega can differ slightly across methods of model identification
  - **Omega is calculated PER FACTOR because it assumes unidimensionality (which should have been tested already)**

# 4 Steps in Model Evaluation: Step 4

- Omega is an estimate of reliability for a sum score (as is alpha)
  - But why would you want to use a sum score after you just built a better measurement model for the latent factor??? That's the entire point of SEM—to estimate relations directly among latent variables!
- If you want to use a factor score instead of a sum score, then you should report **factor score reliability** instead of omega:

- Factor score reliability =  $\frac{\sigma_F^2}{\sigma_F^2 + SE_{FS}^2}$

$\sigma_F^2$ = factor variance from model solution $SE_{FS}^2$ = error variance of EAP factor scores denominator = total variance = T/(T+E)
---

- 
- Either way, if you must simplify your model, you can use the sum score or a factor score as a single indicator of a latent variable, but correct for its unreliability as follows (example using Mplus syntax):
    - `Factor BY fscore@1; fscore* (ResVar); Factor*;  
[fscore*]; [Factor@0]; ! To mean-center latent factor`
    - `MODEL CONSTRAINT: Resvar = (1 - reliability) * (total variance);  
! Factor score residual variance = "unreliable" amount of total variance`
    - **Example using R lavaan syntax:**  
`Factor =~ 1*fscore; fscore ~~ ResVar*fscore; Factor ~~ Factor;  
fscore ~ 1; Factor ~ 0; # Mean center latent factor  
ResVar := (1-reliability)*(total variance)`

# CFA Model Evaluation: Summary

- The primary advantage of working in a CFA framework is obtaining indices of global and local model fit
  - $\chi^2$  and model fit indices indicate how well the model recreates the item means, variances, and covariances relative to the original data...
    - .. But normalized residuals should still be examined for evidence of local misfit (e.g., badly recreated covariances between certain items)
  - Nested model comparisons via rescaled  $-2\Delta LL$  can be conducted in order to compare the fit of augmented or simplified models...
    - ... But don't rely blindly on modification indices (cheat codes) to do so
  - Effect size and significance of model parameters matters, too
    - ... How well are your latent factors really defined anyway? Effect size!
    - Watch out for out-of-bound estimates—this means something is wrong
    - Watch for unreasonable predicted responses—this means you shouldn't be using a linear slope CFA model (so you need a nonlinear slope model)

# Testing CTT Assumptions in CFA

- **Alpha sum-score reliability** assuming three things:
  - The items measure a single, unidimensional latent factor
  - All factor loadings (discriminations) are equal, or that items are “true-score equivalent” or “**tau-equivalent**”
  - **Local independence** (errors are uncorrelated)
- After assessing unidimensionality of each latent factor, we can then test the assumption of **tau-equivalence** via a  $-2\Delta LL$  comparison against a model in which the factor loadings are constrained to be equal
  - If model fit gets worse, the loadings are not equal; items differ in discrimination
  - If so, don't use alpha—use model-based reliability (omega) instead, because omega assumes unidimensionality, but not tau-equivalence
- The assumption of **parallel items** is then testable by constraining item error variances to be equal, too—does model fit get worse?
  - Parallel items is needed to use Spearman-Brown formulas to predict reliability
  - Parallel items will hardly ever hold in real data → **sum scores ≠ factor scores!**
  - Note that if tau-equivalence doesn't hold, then neither does parallel items



# Conclusion: The Big Picture of CFA

- **The CFA unit of analysis is the ITEM:**  $y_{is} = \mu_i + \lambda_i F_s + e_{is}$ 
  - **Linear** regression relating **continuous** item responses to latent factor predictor
  - Both item AND subject properties matter in predicting item responses
  - Latent factors are estimated as separate entities based on the observed covariances among items—latent factors represent testable assumptions
    - Items are unrelated after controlling for factor(s) → local independence
    - Extra item relations can be modeled via error covariances and method factors
- **Because item responses are included:**
  - Items are allowed to vary in discrimination (as factor loadings)  
→ thus, exchangeability (tau-equivalence) is a testable hypothesis
  - Because difficulty (item intercepts) do not contribute to the covariance, they don't really matter in CFA (unless you are testing factor mean differences)
  - To make a test better, you need more items, but not just any kind!
    - **What kind of items? Ones with *higher standardized loadings* (more information)**
  - Measurement error is still assumed constant across the latent trait
    - **People low-medium-high in Factor Score are measured equally well**